

CIPROFLOXACIN (CIPRO®) INDICATIONS

Ciprofloxacin is indicated for the treatment of the following infections:

INFECTION

CAUSE

Acute sinusitis

Haemophilus influenzae
Streptococcus pneumoniae
Moraxella catarrhalis

Lower respiratory tract infections

Escherichia coli
Klebsiella pneumoniae
Enterobacter cloacae
Proteus mirabilis
Pseudomonas aeruginosa
Haemophilus influenzae
Haemophilus parainfluenzae
Streptococcus pneumoniae

Note: Although effective in clinical trials, ciprofloxacin is not a drug of first choice in the treatment of presumed or confirmed pneumonia secondary to *Streptococcus pneumoniae* infection.

Acute exacerbations of chronic bronchitis

Moraxella catarrhalis

Urinary tract infections

Escherichia coli
Klebsiella pneumoniae
Serratia marcescens
Enterobacter cloacae
Proteus mirabilis
Pseudomonas aeruginosa
Providencia rettgeri
Morganella morganii
Citrobacter diversus
Citrobacter freundii
Staphylococcus epidermis
Staphylococcus saprophyticus
Enterococcus faecalis

Acute uncomplicated cystitis in females

Escherichia coli
Staphylococcus saprophyticus

CIPROFLOXACIN (CIPRO®) INDICATIONS

Ciprofloxacin is indicated for the treatment of the following infections:

infection	cause
Chronic bacterial prostatitis	Escherichia coli Proteus mirabilis
Complicated intraabdominal infections (used in combination with metronidazole - Flagyl®; oral, IV)	Escherichia coli Proteus mirabilis Pseudomonas aeruginosa Klebsiella pneumoniae Bacteroides fragilis
Skin and skin structure infections	Escherichia coli Klebsiella pneumoniae Enterobacter cloacae Proteus mirabilis Pseudomonas aeruginosa Proteus vulgaris Providencia stuartii Morganella morganii Citrobacter freundii Staphylococcus aureus (methicillin susceptible) Staphylococcus epidermis Streptococcus pyogenes
Bone and joint infections	Enterobacter cloacae Serratia marcescens Pseudomonas aeruginosa
Infectious diarrhea (when antibacterial therapy is indicated)	Escherichia coli (enterotoxigenic strains) Campylobacter jejuni Shigella boydii Shigella dysenteriae Shigella flexneri Shigella sonnei
Typhoid fever (enteric fever)	Salmonella typhi

CIPROFLOXACIN (CIPRO®) INDICATIONS

Ciprofloxacin is indicated for the treatment of the following infections:

INFECTION

CAUSE

Uncomplicated cervical
and urethral gonorrhoea

Neisseria gonorrhoeae

Nosocomial pneumonia

Haemophilus influenzae
Klebsiella pneumoniae

Empirical therapy for febrile
neutropenic patients

N/A

(used in combination with
piperacillin sodium - Pipracil®)

